

(19) World Intellectual Property Organization
International Bureau



10 Dec 2000 PCT/PTO 13 DEC 2000

(43) International Publication Date
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number
WO 00/77199 A1

(51) International Patent Classification⁷: C12N 15/12,
5/10, C07K 14/705, 16/28, A61K 38/17, G01N 33/50

(21) International Application Number: PCT/EP00/05592

(22) International Filing Date: 14 June 2000 (14.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9913850.5 14 June 1999 (14.06.1999) GB

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turn-
houtseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BENDER, Eckhard
[DE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg
30, B-2340 Beerse (BE). PINDON, Armelle, Nathalie
[FR/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg
30, B-2340 Beerse (BE). VAN OERS, Irma, Petronella
[NL/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg
30, B-2340 Beerse (BE). JURZAK, Mirek [DE/BE];
Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340
Beerse (BE). LUYTEN, Walter, Herman, Maria, Louis
[BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30,
B-2340 Beerse (BE).

(74) Agent: BALDOCK, Sharon, Claire; Boulton Wade Ten-
nant, Verulam Gardens, 70 Gray's Inn Road, London
WC1X 8BT (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: CLONING AND EXPRESSION OF A NOVEL 5-HT₄ RECEPTOR

(57) Abstract: There is disclosed an isolated or substantially pure form of a nucleic acid molecule encoding a human 5-HT_{4(b)} receptor, which 5-HT_{4(b)} receptor preferably comprises the sequence illustrated in Figure 1b. Also provided by the invention are expression vectors incorporating said nucleic acid molecule in addition to transgenic cells, tissues or rganisms transfected with the nucleic acid molecule.



WO 00/77199 A1

CLONING AND EXPRESSION OF A NOVEL 5-HT₄ RECEPTOR

The present invention is concerned with cloning and expression of a novel receptor and, in particular, with a novel nucleic acid sequence encoding a 5-HT₄ receptor splice variant designated herein as 5-HT_{4(h)}, an expression vector comprising said nucleic acid sequence, a host cell transformed or transfected with said vector, the 5-HT_{4(h)} receptor protein expressed from said host cell and pharmaceutical compositions comprising said expressed protein or said nucleic acid or its complementary sequences.

The 5-HT₄ receptor is widely distributed in the body, in the periphery as well as in the central nervous system. In the periphery it is found in the gastrointestinal tract, for example in the esophagus (Moumami et al., 1992), the ileum (Buchheit and Buhl, 1991) and colon (Elswood et al., 1991). It is also present in the atrium (Kaumann et al., 1990), the bladder (Candura et al., 1996) and the adrenal glands. In the rat brain, 5-HT₄ mRNA has been discovered by in situ hybridization in the olfactory tubercle, the striatum and the hippocampus (Vilaro et al., 1996). The wide distribution in different tissues of the 5-HT₄ receptor is paralleled by a wide variety of 5-HT₄ variants caused by alternative splicing of exons. The splice variants described so far (Gerald et al., 1995; Claeysen et al., 1996; Van den Wyngaert et al., 1997; Claeysen et al., 1997; Blondel et al., 1997; Blondel et al., 1998) are all variations of the cytoplasmic C-terminus.

- 2 -

The predicted protein structures encoded by cDNA sequences already known reveal seven transmembrane domains for the complete open reading frames. In addition to their structure and 5-HT₄ receptor coupled signal transduction events (increase in cAMP formation, opening of K⁺ channels), 5-HT₄ receptors have also been classified as G-protein coupled receptors (GPCRs).

10 The present inventors have identified a novel human 5-HT₄ splice variant (h), which leads to the insertion of 14 amino acids into the second extracellular loop of the receptor protein. They expressed the isolated full length cDNA transiently in mammalian cells in order to compare its pharmacology with already known 5-HT₄ splice variants and its tissue distribution is analyzed by RT-PCR.

20 Therefore, according to a first aspect of the present invention, there is provided, an isolated substantially pure form of nucleic acid molecule encoding a human 5-HT_{4(h)} receptor. Preferably the 5-HT_{4(h)} receptor encoded by said nucleic acid molecule comprises the amino acid sequence illustrated in Figure 1b or a functional equivalent, derivative or bioprecursor of said receptor.

Thus, the present invention comprises a nucleic acid molecule encoding a human 5-HT_{4(h)} receptor or an immunologically and/or biologically active fragment thereof, which comprises a nucleotide sequence selected from the group consisting of:

(a) nucleotide sequences encoding the amino acid

- 3 -

sequence depicted in Figure 1b;

- (b) nucleotide sequences comprising the coding sequence as depicted in Figure 1a;
- (c) nucleotide sequences encoding a polypeptide
5 derived from the polypeptide encoded by a nucleotide sequence of (a) or (b) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence encoded by the nucleotide sequence
10 of (a) or (b);
- (d) nucleotide sequences the complementary strand of which hybridises with a nucleotide sequence of any one of (a) to (c);
- (e) nucleotide sequences encoding a polypeptide
15 the amino acid sequence of which has an identity of 70% or more to the amino acid sequence of the polypeptide encoded by a nucleotide sequence of any one of (a) to (d);
- (f) nucleotide sequences encoding a polypeptide
20 capable of binding a ligand of 5-HT_{4(h)} comprising a fragment or an epitope-bearing portion of a polypeptide encoded by a nucleotide sequence of any one of (a) to
25 (e);
- (g) nucleotide sequences comprising at least 15 consecutive nucleotides of a nucleotide sequence of any one of (a) to (f);
- (h) nucleotide sequences comprising a nucleotide
30 sequence which is degenerated as a result of the genetic code to a nucleotide sequence of any of (a) to (g).

- 4 -

Advantageously, the isolated nucleic acid according to the invention may be used for expression in, for example, a host cell or the like using a suitable expression vector. Preferably, the nucleic acid may
5 be a DNA molecule or a cDNA molecule. Preferably, the DNA molecule has the nucleic acid sequence as illustrated in Figure 1a.

The nucleic acid molecule is preferably capable of
10 hybridising to the sequences of the invention under conditions of high stringency or to the complement thereof.

Stringency of hybridisation as used herein refers to
15 conditions under which polynucleic acids are stable. The stability of hybrids is reflected in the melting temperature (T_m) of the hybrids. T_m can be approximated by the formula:

$$20 \quad 81.5^{\circ}\text{C} + 16.6(\log_{10}[\text{Na}^+] + 0.41 (\% \text{G\&C}) - 6001/l$$

wherein l is the length of the hybrids in nucleotides. T_m decreases approximately by 1-1.5°C with every 1% decrease in sequence homology.

25 The term "stringency" refers to the hybridisation conditions wherein a single-stranded nucleic acid joins with a complementary strand when the purine or pyrimidine bases therein pair with their corresponding
30 base by hydrogen bonding. High stringency conditions favour homologous base pairing whereas low stringency conditions favour non-homologous base pairing.

- 5 -

"Low stringency" conditions comprise, for example, a temperature of about 37°C or less, a formamide concentration of less than about 50%, and a moderate to low salt (SSC) concentration; or, alternatively, a
5 temperature of about 50°C or less, and a moderate to high salt (SSPE) concentration, for example 1M NaCl.

"High stringency" conditions comprise, for example, a temperature of about 42°C or less, a formamide
10 concentration of less than about 20%, and a low salt (SSC) concentration; or, alternatively, a temperature of about 65°C, or less, and a low salt (SSPE) concentration. For example, high stringency conditions comprise hybridization in 0.5 M NaHPO₄, 7%
15 sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C (Ausubel, F.M. et al. Current Protocols in Molecular Biology, Vol. I, 1989; Green Inc. New York, at 2.10.3).

20 "SSC" comprises a hybridization and wash solution. A stock 20X SSC solution contains 3M sodium chloride, 0.3M sodium citrate, pH 7.0.

"SSPE" comprises a hybridization and wash solution. A
25 1X SSPE solution contains 180 mM NaCl, 9mM Na₂HPO₄ and 1 mM EDTA, pH 7.4.

The nucleic acid capable of hybridising to nucleic acid molecules according to the invention will
30 generally be at least 70%, preferably at least 80 or 90% and more preferably at least 95% homologous to the nucleotide sequences according to the invention.

- 6 -

Advantageously, the antisense molecule may be used as a probe or as a medicament or in a pharmaceutical composition together with a pharmaceutically acceptable carrier, diluent or excipient.

5

The term "homologous" describes the relationship between different nucleic acid molecules or amino acid sequences wherein said sequences or molecules are related by partial identity or similarity at one or more blocks or regions within said molecules or sequences. Homology may be determined by means of computer programs known in the art.

Substantial homology preferably carries with it that the nucleotide and amino acid sequences of the 5-HT_{4(h)} of the invention comprise a nucleotide and amino acid sequence fragment, respectively, corresponding and displaying a certain degree of sequence identity to the sequences in Figure 1a and 1b. Preferably they share an identity of at least 30 %, preferably 40 %, more preferably 50 %, still more preferably 60 %, most preferably 70%, and particularly an identity of at least 80 %, preferably more than 90 % and still more preferably more than 95 % is desired with respect to the nucleotide or amino acid sequences depicted in Figures 1a and 1b, respectively. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using, for example, the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6 (1990), 237-245.) In a sequence alignment the query

- 7 -

and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Further programs that can be used in order to determine homology/identity are described below and in the examples. The sequences that are homologous to the sequences described above are, for example, variations of said sequences which represent modifications having the same biological function, in particular encoding proteins with the same or substantially the same receptor specificity, e.g. binding specificity. They may be naturally occurring variations, such as sequences from other mammals, or mutations. These mutations may occur naturally or may be obtained by mutagenesis techniques. The allelic variations may be naturally occurring allelic variants as well as synthetically produced or genetically engineered variants. In a preferred embodiment the sequences are derived from a human.

According to a further aspect of the invention, there is provided a DNA expression vector comprising the DNA molecule according to the invention. This vector may advantageously be used to transform or transfect a host cell to achieve expression of the 5-HT_{4(h)} receptor from said cell. Preferably, the DNA molecule is included in a plasmid such as, for example, pcDNA3 for subsequent transformation or transfection of said host cell.

An expression vector according to the invention includes a vector having a nucleic acid according to the invention operably linked to regulatory sequences,

- 8 -

such as promoter regions, that are capable of effecting expression of said DNA fragments. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship
5 permitting them to function in their intended manner. Such vectors may be transformed into a suitable host cell to provide for expression of a polypeptide according to the invention. Thus, in a further aspect, the invention provides a process for preparing
10 polypeptides according to the invention which comprises cultivating a host cell, transformed or transfected with an expression vector as described above under conditions to provide for expression by the vector of a coding sequence encoding the
15 polypeptides, and recovering the expressed polypeptides.

The vectors may be, for example, plasmid, virus or phage vectors provided with an origin of replication,
20 optionally a promoter for the expression of said nucleotide and optionally a regulator of the promoter. The vectors may include a coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the invention. In
25 addition the vector may contain a sequence coding a phenotypic trait for selection of transformed cells such as, for example, ampicillin resistance.

Regulatory elements required for expression include
30 promoter sequences to bind RNA polymerase and transcription initiation sequences for ribosome binding. For example, a bacterial expression vector may include a promoter such as the lac promoter and

- 9 -

for transcription initiation in the Shine-Dalgarno sequence and the start codon AUG. Similarly, a eukaryotic expression vector may include a heterologous or homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome. Such vectors may be obtained commercially or assembled from the sequences described by methods well known in the art.

10

Transcription of DNA encoding the polypeptides of the present invention by higher eukaryotes is optimised by including an enhancer sequence in the vector. Enhancers are cis-acting elements of DNA that act on a promoter to increase the level of transcription. Vectors will also generally include origins of replication in addition to the selectable markers.

15

20

Nucleic acid molecules according to the invention may be inserted into the vectors described in an antisense orientation in order to provide for the production of antisense RNA. Antisense RNA or other antisense nucleic acids may be produced by synthetic means.

25

30

In accordance with the present invention, a defined nucleic acid includes not only the identical nucleic acid but also any amino base variations including, in particular, substitutions in bases which result in a synonymous codon (a different codon specifying the same amino acid residue) due to the degenerate code in conservative amino acid substitutions. The term "nucleic acid sequence" also includes the complementary sequence to any single stranded sequence

- 10 -

given regarding base variations.

The present invention also advantageously provides nucleic acid sequences of at least approximately 10
5 contiguous nucleotides of a nucleic acid according to the invention and preferably from 10 to 50 nucleotides. These sequences may, advantageously, be used as probes or primers to initiate replication, or the like. Such nucleic acid sequences may be produced
10 according to techniques well known in the art, such as, by recombinant or synthetic means. They may also be used in diagnostic kits or the like for detecting the presence of a nucleic acid according to the invention. These tests generally comprise contacting
15 the probe with the sample under hybridising conditions and detecting for the presence of any duplex or triplex formation between the probe and any nucleic acid in the sample.

20 According to the present invention these probes may be anchored to a solid support. Preferably, they are present on an array so that multiple probes can simultaneously hybridize to a single biological sample. The probes can be spotted onto the array or
25 synthesised *in situ* on the array. (See Lockhart et al., Nature Biotechnology, vol. 14, December 1996 "Expression monitoring by hybridisation into high density oligonucleotide arrays". A single array can contain more than 100, 500 or even 1,000 different
30 probes in discrete locations.

The nucleic acid sequences, according to the invention may be produced using such recombinant or synthetic

- 11 -

means, such as, for example, using PCR cloning mechanisms which generally involve making a pair of primers, which may be from approximately 10 to 50 nucleotides to a region of the gene which is desired to be cloned, bringing the primers into contact with mRNA, cDNA, or genomic DNA from a human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified region or fragment and recovering the amplified DNA. Generally, such techniques as defined herein are well known in the art, such as described in Sambrook et al (Molecular Cloning: a Laboratory Manual, 1989).

The nucleic acids or oligonucleotides according to the invention may carry a revealing label. Suitable labels include radioisotopes such as ^{32}P or ^{35}S , enzyme labels or other protein labels such as biotin or fluorescent markers. Such labels may be added to the nucleic acids or oligonucleotides of the invention and may be detected using known techniques *per se*.

Advantageously, human allelic variants or polymorphisms of the DNA molecule according to the invention may be identified by, for example, probing cDNA or genomic libraries from a range of individuals, for example, from different populations. Furthermore, nucleic acids and probes according to the invention may be used to sequence genomic DNA from patients using techniques well known in the art, such as the Sanger Dideoxy chain termination method, which may, advantageously, ascertain any predisposition of a patient to certain disorders associated with a growth

- 12 -

factor according to the invention.

The present invention also comprises within its scope proteins or polypeptides encoded by the nucleic acid molecules according to the invention or a functional equivalent, derivative or bioprecursor thereof. Preferably, the protein comprises the amino acid sequence illustrated in Figure 1b.

10 A "functional equivalent" as defined herein should be taken to mean a receptor that exhibits the same properties and functionality associated with the 5HT_{4(h)} receptor according to the invention. A "derivative" should be taken to mean a polypeptide or protein in
15 which certain amino acids may have been altered or deleted or replaced and which polypeptide or protein retains biological activity of said 5HT_{4(h)} receptor and/or which can cross react with antibodies raised using a receptor according to the invention as the
20 challenging antigen.

Encompassed with the scope of the invention are hybrid and modified forms of the 5HT_{4(h)} receptor according to the invention including fusion proteins and fragments.
25 The hybrid and modified forms include, for example, when certain amino acids have been subjected to some modification or replacement, such as for example, by point mutation and yet which results in a protein which possesses the same receptor specificity as the
30 5HT_{4(h)} of the invention.

The protein according to the invention should be taken to include all possible amino acid variants encoded by

- 13 -

the nucleic acid molecule according to the invention including a polypeptide encoded by said molecule and having conservative amino acid changes. Proteins or polypeptides according to the invention further
5 include variants of such sequences, including naturally occurring allelic variants which are substantially homologous to said proteins or polypeptides. In this context, substantial homology is regarded as a sequence which has at least 70%, and
10 preferably 80 or 90% amino acid homology with the proteins or polypeptides encoded by the nucleic acid molecules according to the invention.

As is well known in the art many proteins are produced
15 *in vivo* with a (pre) signal at the N terminus of the protein and which may be required for transport of the protein across the cell membrane. Furthermore, such proteins may comprise a further pro sequence that represents a stable precursor to the mature protein.
20 Such pre and pro sequences are not required for biological activity. Furthermore, in eukaryotic organisms many proteins are subjected to glycosylation so as to confer biological activity *in vivo*.
References to a bioprecursor, in accordance with the
25 present invention, refers to all such forms of the protein or polypeptide of the invention prior to any such post translational modification.

A further aspect of the invention comprises the host
30 cell itself transformed with the DNA expression vector described herein, which host cell preferably comprises a eukaryotic cell, which may be for example, a mammalian cell, an insect cell or yeast cell or the

- 14 -

like. In one embodiment the cell comprises a human embryonic kidney cell and preferably a cell of the HEK293 cell line. Alternatively, the cell may comprise NIH/3T3 mouse fibroblasts or Chinese hamster ovary (CHO) cells or COS-7 cells.

Further provided by the present invention is a transgenic cell, tissue or organism comprising a transgene capable of expressing a human 5-HT_{4(h)} receptor according to the invention, or expressing a functional equivalent, fragment, derivative or bioprecursor of said receptor. The term "transgene capable of expression" as used herein means a suitable nucleic acid sequence which leads to the expression of a human 5-HT_{4(h)} receptor having the same function and/or activity. The transgene may include, for example, genomic nucleic acid isolated from human cells or synthetic nucleic acid including cDNA integrated into the genome or in an extra chromosomal state. Preferably, the transgene comprises the nucleic acid sequence encoding the 5-HT_{4(h)} receptor as described above, or a functional fragment of said nucleic acid. A functional fragment of said nucleic acid should be taken to mean a fragment of the gene comprising said nucleic acid, coding for the 5-HT_{4(h)} receptor or a functional equivalent, derivative or bioprecursor of said receptor. For example, the gene may comprise deletions or mutations but may still encode a functional 5-HT_{4(h)} receptor protein.

30

There is also provided by a further aspect of the present invention, a purified human 5-HT_{4(h)} receptor expressed by a host cell or a transgenic cell tissue

- 15 -

or organism according to the invention. Also provided by the invention are membrane preparations from cells expressing a human 5-HT_{4(h)} receptor.

5 The nucleic acid or protein according to the invention may be used as a medicament or in the preparation of a medicament for treating cancer or other diseases or conditions associated with expression of 5-HT_{4(h)} receptor protein.

10

Advantageously, the nucleic acid molecule or the protein according to the invention may be provided in a pharmaceutical composition together with a pharmacologically acceptable carrier, diluent or
15 excipient therefor.

The present invention is further directed to inhibiting 5-HT_{4(h)} *in vivo* by the use of antisense technology. Antisense technology can be used to
20 control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the mature protein sequence, which encodes for the protein of the
25 present invention, is used to design an antisense RNA oligonucleotide of from 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple-helix - see Lee et al. Nucl. Acids Res.,
30 6:3073 (1979); Cooney et al., Science, 241:456 (1988); and Dervan et al., Science, 251: 1360 (1991), thereby preventing transcription and the production of 5-HT_{4(h)}. The antisense RNA oligonucleotide hybridises to the

- 16 -

mRNA *in vivo* and blocks translation of an mRNA molecule into the 5-HT_{4(h)} receptor.

5 Alternatively, the oligonucleotide described above can be delivered to cells by procedures in the art such that the anti-sense RNA and DNA may be expressed *in vivo* to inhibit production of a polypeptide of the invention in the manner described above.

10 Antisense constructs to the nucleotide sequence encoding 5-HT_{4(h)}, therefore, may inhibit the expression of the 5-HT_{4(h)} receptor and may therefore be used to treat conditions associated with expression or overexpression of 5-HT_{4(h)} according to the invention.

15 A further aspect of the invention comprises the host cell itself transformed with the DNA expression vector described herein, which host cell preferably comprises a mammalian cell such as, for example, a COS-7 cell or
20 a human cell such as a human embryonic kidney (HEK) 293 cell or the like.

Incorporation of cloned DNA into a suitable expression vector for subsequent transformation of the cell and
25 subsequent selection of the transformed cells is well known to those skilled in the art as provided in Sambrook et al., (1989) "Molecular Cloning, A Laboratory Manual, Cold Spring Harbour Laboratory Press".

30 Antibodies to the human 5-HT_{4(h)} receptor are also provided which may be used in a medicament or in a pharmaceutical composition.

- 17 -

Antibodies to the protein or polypeptide of the present invention may, advantageously, be prepared by techniques which are known in the art. For example, polyclonal antibodies may be prepared by inoculating a host animal, such as a mouse, with the polypeptide according to the invention or an epitope thereof and recovering immune serum. Monoclonal antibodies may be prepared according to known techniques such as described by Kohler R. and Milstein C., Nature (1975) 256, 495-497.

Antibodies according to the invention may also be used in a method of detecting for the presence of a receptor according to the invention, which method comprises reacting the antibody with a sample and identifying any protein bound to said antibody. A kit may also be provided for performing said method which comprises an antibody according to the invention and means for reacting the antibody with said sample.

Advantageously, the antibody according to the invention may also be used as a medicament or in the preparation of a medicament for treating diseases associated with expression of 5-HT_{4(h)}. The invention also further provides a pharmaceutical composition comprising said antibody together with a pharmaceutically acceptable carrier, diluent or excipient therefor.

Proteins which interact with the polypeptide of the invention may be identified by investigating protein-protein interactions using the two-hybrid vector system first proposed by Chien et al (1991), Proc.

Natl. Acad. Sci. USA 88 : 9578-9582.

This technique is based on functional reconstitution
in vivo of a transcription factor which activates a
5 reporter gene. More particularly the technique
comprises providing an appropriate host cell with a
DNA construct comprising a reporter gene under the
control of a promoter regulated by a transcription
factor having a DNA binding domain and an activating
10 domain, expressing in the host cell a first hybrid DNA
sequence encoding a first fusion of a fragment or all
of a nucleic acid sequence according to the invention
and either said DNA binding domain or said activating
domain of the transcription factor, expressing in the
15 host at least one second hybrid DNA sequence, such as,
a library or the like, encoding putative binding
proteins to be investigated together with the DNA
binding or activating domain of the transcription
factor which is not incorporated in the first fusion;
20 detecting any binding of the proteins to be
investigated with a protein according to the invention
by detecting for the presence of any reporter gene
product in the host cell; optionally isolating second
hybrid DNA sequences encoding the binding protein.

25 Proteins which bind to the 5-HT_{4(h)} receptor can be
identified using this technique. The proteins
identified can also be used to identify compounds
which acts as agonists/antagonists of these proteins.
30 The structure of the receptor can also be used to
design agonists or antagonists of the receptor. The
present invention also comprises an agonist or
antagonist of the human 5-HT_{4(h)} receptor according to

- 19 -

the invention which agonist or antagonist advantageously may also be used as a medicament or in a pharmaceutical composition together with a pharmaceutically acceptable carrier diluent or
5 excipient therefor.

The present invention is also directed to antagonists and inhibitors of the 5-HT_{4(h)} receptor of the present invention. The antagonists and inhibitors are those
10 substances which inhibit or eliminate the function of such a receptor. The present invention further relates to agonists and stimulators of a receptor of the present invention. The agonists and stimulators are those substances which enhance the function or
15 activity or the expression of such a receptor.

Further provided by the present invention is a method of determining whether a compound is an agonist or an antagonist of a 5-HT_{4(h)} receptor protein, which method
20 comprises contacting a host cell or transgenic cell tissue or organism according to the invention expressing said 5-HT_{4(h)} receptor protein with said compound in the presence of a protein which binds to said receptor and monitoring induced cAMP formation in
25 said cell. Preferably, the cell is a mammalian cell such as a COS-7 cell or the like or a human cell, such as a human embryonic kidney (HEK) 293 cell or the like. A further method of determining whether a compound is an agonist or an antagonist of 5-HT_{4(h)}
30 ligand protein is provided which method comprises contacting a cell or membrane preparation of said host cell or said transgenic cell according to the invention with said compound and establishing the

- 20 -

binding affinity of said compound for said receptor.
Any compounds identified may advantageously be used as
a medicament or in a pharmaceutical composition
together with an appropriate diluent or excipient.

5

Antagonists of the 5-HT_{4(h)} receptor according to the
invention may be useful in treating any of heartburn,
reflux, esophagitis, Barrett's esophagus, esophageal
cancer, achalasia, esophageal stenosis, esophageal
10 spasms, esophageal hiatal hernia or other esophageal
motility disorders, oesophageal irritation, such as
asthma, bronchospasms, aspiration and its consequences
(bronchitis, (broncho)pneumonia, bronchiectasia) and
other diseases of the lower oesophageal sphincter, or
15 achalasia; oesophageal stenosis (due to systemic
sclerosis, tumours, burns, or the like) or
compression, oesophageal spasms or other oesophageal
motility disorders, asthma, irritable bowel syndrome,
bronchospasms and other airway disorders possibly
20 connected with oesophageal irritation aspiration and
its consequence (bronchitis, (broncho)pneumonia,
bronchiectasia); (hiatus) hernia; denervation of the
oesophagus (e.g. after certain types of trauma or
surgery), disturbances in oesophageal innervation.

25

Further provided by the present invention is a
diagnostic kit for determining whether a compound is
an agonist or an antagonist in relation to 5-HT_{4(h)}
receptor ligand or an antibody thereto. The kit may
30 comprise a cell according to the invention, means for
contacting said compound with said cell and means for
measuring cAMP formation in said cell. Alternatively,
the kit may comprise a probe, including any of a

- 21 -

nucleic acid molecule encoding a 5-HT_{4(h)} receptor according to the invention, a molecule capable of hybridising thereto under high stringency conditions, a fragment of said nucleic acids, an antisense molecule according to the invention, together with means for contacting biological material to be tested with said probe.

Disorders mediated by activation or expression of the 5-HT_{4(h)} receptor may, advantageously, be treated by administering to an individual an amount of a compound identified as an agonist of the ligand binding 5HT_{4(h)} in sufficient concentration to reduce or prevent the symptoms of the disorder.

The present invention may be more clearly understood from the following exemplary embodiment with reference to the accompanying figures wherein;

Figure 1: a) is an alignment of nucleotide sequences of dog and human 5-HT_{4(h)}. The positions of primers used in this study are indicated by arrows and b) is an alignment of amino acid sequences of dog and human 5-HT_{4(h)}.

Figure 2: is an illustration of mRNA tissue distribution performed as described in Materials and Methods. The letters indicate the used primer combination in the PCR, A:FW AB1/REV B1 (5'part 5-HT₄ cDNA including h exon), B:FW AB1/REV AB2 (common part of all 5-HT₄ splice

- 22 -

5 variants), C:FW AB2/REV SH1 (3'part 5-HT_{4a} cDNA), D:FW AB2/REV LO1 (3'part 5-HT_{4b} cDNA), E:FW B1/REV SH1 (3'part 5-HT₄ cDNA, combination of exon h and a), F:FWB1/REV LO1 (3'part 5-HT_{4(h)}).

10 Figure 3a: Saturation analysis of [³H]GR113808 binding on membrane preparation from COS-7 cells transfected with the h 5-HT_{4(h)}. B. Saturation analysis of [³H]5-HT binding on membrane preparation from COS-7 cells transfected with the h 5-HT_{4(h)}.

15 Figure 4: Inhibition of specific [³H]GR113808 binding by 5-HT₄ agonist and antagonist. Membrane preparations from COS-7 cells transiently transfected with h 5-HT_{4(h)} receptor were incubated with 0.25 nM [³H]GR113808. Non-specific binding was determined by 10 mM SB204070. Results are percentages, 100% is defined by specific binding in the absence of competing compound. Results are the mean of three independent experiments from three different transfections. Calculated pIC₅₀ values are given in Table 1.

25
30 Figure 5: Indirect estimation of AC stimulation by measuring cAMP formation in COS-7 cells transiently transfected with h 5-HT_{4(h)}. Results represent the increase of cAMP after stimulation by agonist since

- 23 -

basal level have been removed. Results are the mean of three independent experiments from three different transfections. Calculated pEC50 and % of 5-HT_{max} values are given in Table 2. The efficacy and potency of the different agonists to trigger the cellular response was estimated and compared for the three different variants. The mean of pEC50 and the percentage of stimulation, normalized for the maximum stimulation induced by 5-HT (% of 5-HT maximum) for the h5-HT_{4(h)}, h5-HT_{4(a)} and h 5-HT_{4(b)}, are presented in Table 2. No difference in the pEC50 was noticed. The cAMP assay has been performed also for COS-7 cells transfected with the empty vector as a negative control. After stimulation with 10⁻⁶ M of each agonist, 5-HT, cisapride and prucalopride, no significant increase of the cAMP basal level was found.

Materials and Methods

Materials

5 AmpliTaq Gold, dNTPs, MgCl₂, and PCR bufferII were
obtained from Perkin-Elmer Cetus (Foster City, CA,
U.S.A.). T4 DNA ligase and restriction endonucleases
were products of Boehringer (Mannheim, Germany). The
Multiprime DNA labeling system and [³H]GR 113808 with
10 a specific activity of 3.07 TBq/mmol were obtained
from Amersham (Little Chalfort, U.K.). The
ExpressHybTM hybridization solution and the Smart cDNA
synthesis kit were from Clontech (Palo Alto, CA,
U.S.A.). [³²P] dCTP was purchased from NEN DuPont
15 (Wilmington, DE, U.S.A.). Plasmid preparation kits and
the Qiaquick PCR amplification kit were from Qiagen
(Hilden, Germany). The PRISM Ready Reaction Dye
Terminator Cycle Sequencing kits and the ABI 377 or
373A sequencing machines were from Applied Biosystems
20 (Foster City, CA, U.S.A.). The Geneamp PCR System 9600
was from Perkin-Elmer (Norwalk, CT, U.S.A.). The
mammalian expression vector pcDNA3 was obtained from
Invitrogen (Carlsbad, CA, U.S.A.). Dulbecco's modified
Eagle medium (DMEM) and foetal calf serum were from
25 Life Technologies (Gaithersburg, MD, U.S.A.). The
Bradford protein assay was performed with the reagent
supplied from Bio-Rad (Nazareth Eke, Belgium), which
also supplied the Zeta-Probe blotting membrane. The
NEN flash plate assay was supplied by DuPont de
30 Nemours (Brussels, Belgium). The liquid scintillation
spectrometer and the scintillation fluid Ultima Gold
MV were from Packard (Meriden, CT, U.S.A.). All
compounds were dissolved and diluted in dimethyl

- 25 -

5 sulfoxide (DMSO; except the indoleamines, which were dissolved in water and protected from light throughout the experiment). The final DMSO concentration in the test did not exceed 0.5% (vol/vol). The GraphPad Prism program was from GraphPad Software, Inc. (San Diego, CA, U.S.A.).

General molecular biological methods

10 Unless otherwise indicated, all PCR reactions were performed in a total volume of 50 ml, containing 1 ml of cDNA and 1.25 U of AmpliTaq Gold in 1x PCR buffer II, 200 mM dNTPs, 400 nM primers, and 2.0 mM MgCl₂.
15 PCR conditions were 10 min of denaturation at 95° C, followed by 35 cycles of 10 seconds at 95° C, 30 seconds at 53° C, and 2 min at 72° C, followed by a 10 min incubation at 72° C. DNA manipulations were done according to standard protocols (Maniatis et al., 1982). DNA sequencing was carried out with reagents
20 from the PRISM Ready Reaction Dye Terminator Kit and run on a GeneAmp PCR System 9600 according to the specifications of the supplier.

BAC library screening

25 A human genomic DNA library in pBeloBAC11, Research Genetics (Huntsville, AL, U.S.A.) was screened by PCR using two primers, FW AB3 5'CTTCATGGTCAACAAGCCCTAC 3' and REV AB2 5'CCCGTTGTAACATCTGGATTGTVYGGGC3', specific
30 for the 5-HT₁ cDNA. The position of the primers on the cDNA sequence is indicated in Fig.1. The PCRs were set up as described above in a total volume of 30 ml, 1 ml of the BAC pools supplied by Research Genetics was

- 26 -

used as substrate. BAC DNA was prepared using the Qiagen Maxi preparation kit (Hilden, Germany).

5 **PCR amplification of the 5' and 3' ends of the human 5-HT_{4(h)} and assembly to a full length coding region**

Based on the nucleotide sequence of the 5-HT_{4(h)} specific exon, derived from BAC clone 228K23 (Research
10 Genetics), 2 primers were designed. Forward primer FW B1 (5'GAAAGGAGTCTAAACCAAGGCCT3') and reverse primer REV B2 (5'CGCATGAAAA TCCTGGCCCAGGCCTTGTT3') hybridizing at positions indicated in Fig. 1A. Primer FW B1 was combined with reverse primer REV 3non
15 (5'CAAGCAGCAGCTTAGGACCTG3') and reverse primer REV B2 was combined with forward primer FW ONstart (5'CCACTC ATGCTTATTTCTGTAATG3'). PCR reactions were set up on cDNA prepared from human lower esophageal sphincter using Advantage Taq and initial denaturation for 1 min
20 at 95° C. The resulting PCR products were cloned into EcoRV cut and dephosphorylated pcDNA3 (Invitrogen, Carlsbad, CA, U.S.A.). PCR products representing 5' and 3' part of the 5-HT_{4(h)} were blunted by Klenow treatment and subsequently digested with the
25 restriction enzyme StyI before ligation into pcDNA3.

mRNA tissue distribution analysis

Total RNA from the different tissues analysed was
30 prepared by the CsCl method, cDNA was prepared thereof using the Smart cDNA library kit from Clontech (Palo Alto, CA, U.S.A.), 0.5 ml of the reaction product was used per PCR. The tissue distribution experiments were

- 27 -

done by PCRs, using 3 different forward primers and 4 different reverse primers. One forward primer, FW B1 is specific for cDNAs containing the 5-HT_{4(h)} exon, the two others FW AB1 (5'GRAAYAAGATGACCCCTCTRCGYATC3') and FW AB2 (5'GCCCRNCARATCCAGATGTTACAACG3') will amplify all 5-HT₄ messages. Among the four reverse primers, one REV AB2 will amplify all 5-HT₄ messages, the other three reverse primers are specific for 5-HT_{4(a)} (REV SH1, 5'GTATGGGCARYTTCTCS AGT TCCTGRTGWTG3'), 5-HT_{4(b)} (REV LO1, 5'GAASTTGCTGNVRGGTGRCACYGACTCTC3') and 5-HT_{4(h)} (REV B2). The position of the primers is indicated in figure 1. The PCR reactions were separated on an agarose gel and blotted on Zeta-Probe blotting membrane. A ³²P-labeled probe (Multiprobe DNA labeling system (Amersham)) corresponding to the human 5-HT_{4(b)} cDNA (Van den Wyngaert et al., 1997) was hybridized to the PCR products on the membrane according to the instructions of the ExpressHybTM user manual (Clontech, Palo Alto, CA, U.S.A.).

Expression of the human 5-HT_{4(h)} receptor in mammalian cells and pharmacological characterization

COS-7 cells were grown in DMEM supplemented with 10% fetal calf serum. A large scale plasmid preparation of 5-HT_{4(h)}/pcDNA3 was made using the Qiagen large scale plasmid preparation kit. Plasmid DNA was transfected into COS-7 cells as described in Van den Wyngaert et al. (1997). 48 hours after transfection the cells were harvested and used for membrane preparation or cyclic AMP formation assays as described (Van den Wyngaert et al., 1997).

- 28 -

Membrane preparation

The transfected COS-7 cells were cultured on 150 mm Petri dishes and washed twice with ice-cold phosphate-buffered saline. The cells were then scrapped from the plates with a cell scraper, suspended in 50 mM Tris-HCl buffer, pH 7.4, and harvested by centrifugation for 10 min at 16000 g. The pellet was resuspended in 5 mM Tris-HCl, pH 7.4, and homogenized with an Ultra Turax homogenizer; the resulting membranes were collected by centrifugation for 20 min at 25000g. Membranes were stored at -70° C in 50 mM Tris-HCl buffer , pH 7.4, at a protein concentration of 1 mg/ml. The Bradford protein assay was used for protein determination with bovine serum albumin as a standard.

Radioligand binding

Assay mixtures (0.5 ml) contained 50 μ l of the tritiated ligand, (either the 5-HT₄ antagonist [³H]GR113808, or the agonist [³H]5-HT), 0.4 ml of membrane preparation (at 0.012 mg/ml of protein for [³H]GR113808 binding or 0.1mg/ml for [³H]5-HT), and 50 μ l solvent for total binding, or 50 μ l of 10 mM SB204070 to determine non specific binding. The [³H]GR113808 assay buffer was 50 mM HEPES/NaOH pH 7.5. The [³H]5-HT assay buffer was Tris-HCl pH 7.4 containing 10 mM MgCl₂, 1 mM pargyline (monoamine oxidase inhibitor) and 1 mM paroxetine (5-HT transport inhibitor). The mixture was incubated 1 hour at 25 °C. The incubation was terminated by rapid filtration over Whatman GF/B filters presoaked in 0.15% polyethylenimine and three washing steps with 3 ml of

- 29 -

50 mM HEPES/NaOH pH 7.5 for [³H]GR113808 binding, presoaked and three washing steps with 3 ml Tris-HCl pH 7.4 for [³H]5-HT binding. Ligand concentration isotherms were obtained using 8 concentrations of [³H]GR113808 from 20 pM to 0.8 nM, and for [³H]5-HT either 8 concentrations from 0.2 nM to 6 nM or from 0.2 nM to 40 nM were chosen. Competition binding experiments were performed with 0.25 nM of [³H]GR113808 and otherwise at the same conditions as for [³H]GR113808 saturation binding. Ligand concentration binding isotherms (rectangular hyperbola) and sigmoidal inhibition curves were calculated by nonlinear regression analysis according to algorithms described by Oestreicher and Pinto (1987). The maximal number of binding sites (B_{max}) and equilibrium dissociation constant (K_D) of the radioligand and the pIC₅₀ (negative logarithm of the concentration that inhibits 50% of specific binding by the radioligand) values of competitors were derived from the curve fitting. Apparent inhibition constant (K_i) values were calculated according to the equation of Cheng and Prusoff (1973). Graphs were prepared using the GraphPad Prism program.

25 **Measurement of cAMP formation**

These experiments were done using the NEN adenylyl cyclase activation flashplate assay, according to the supplier. Cells were removed from the Petri dishes with 3 ml EDTA (0.04% w/v) and resuspended with phosphate buffered saline without Ca²⁺ and Mg²⁺. The cells were centrifuged at 1500 g for 5 minutes and the supernatants were removed. The pellet was resuspend in

- 30 -

stimulation buffer and diluted to a concentration of 10^6 cells/ml, 50 μ l thereof were added per well of the flashplate (50000 cells/well). Compounds were diluted in PBS containing 1 mM pargyline and 1 mM paroxetine, and 50 μ l of the resulting mixture was added per well, followed by an incubation for 20 minutes at 37° C. The final concentration of DMSO (whenever needed to dissolve the compounds) did not exceed 0.5% (vol/vol) and was also included in the corresponding control samples. The experiment was stopped and a direct cAMP [125 I] detection assay was performed by adding 100 μ l of detection mix per well. After incubation for 24h at room temperature, counting was done in a Topcount (Packard).

Results

Cloning of the human 5-HT_{4(h)} splice variant

In the course of cloning the canine 5-HT₄ receptor cDNA by degenerate primer PCR based on our human 5-HT_{4(b)} sequence (Van den Wyngaert et al., 1997) we found a variant (Fig. 1), similar to the partial porcine 5-HT_{4(h)} cDNA originally designated 5-HT₄B (published by Ullmer et al., 1995). In order to amplify this sequence also from human cDNA, we applied primers based on the d 5-HT_{4(h)} specific sequence to different human cDNAs. However we succeeded only in amplifying the 5' part of a putative human homologue and did not receive PCR products for the 3' part of the ORF. In order to investigate, whether there is an extensive sequence diversity between the human and canine version of this exon, respectively whether this exon

- 31 -

exists at all in human, we screened a human genomic DNA BAC library for a clone containing the human 5-HT₄ gene. One positive clone, 228K23 (Research Genetics) was identified. DNA prepared from this clone was

5 sequenced using primer FW B2 (5'AACCAAGGCCTGGGCCAGGATTTTC ATGGG3'), complementary to a part of the 5-HT_{4(h)} exon. The resulting sequence stretched into the adjacent intron sequence, design of a reverse primer complementary to that intron sequence

10 allowed complete sequence determination of the human 5-HT_{4(h)} exon sequence. Based on this information, h-variant specific forward and reverse primers were designed and combined with primer FW ONstart and reverse primers specific for the a and b splice

15 variant. However for the latter, only the b variant specific primer REV 3non produced a PCR product of expected size in combination with FW B1. The two PCR products were fused by using the unique StyI restriction enzyme site in the nucleotide sequence of

20 the 5-HT_{4(h)} specific exon to build a full length reading frame.

Tissue distribution of the 5-HT_{4(h)} mRNA

25 In an initial experiment to explore the specific function of the 5-HT_{4(h)} in human physiology, we performed a tissue distribution study. The primers were chosen in order to obtain PCR products from parts of the 5-HT₄ cDNA that are common to all different

30 variants and also to obtain bands that are specific for the a, b or h exon (Fig.2). The only tissue from which detectable levels of a PCR product corresponding to the 5-HT_{4(h)} variant, could be produced, was the

- 32 -

lower esophageal sphincter (LES). For a number of the examined tissues, we found PCR products corresponding to the presence of h exon mRNA, but not in combination with the a or b specific C-terminal exon, in these cases, the h specific exon may be fused to another C-terminal exon of the 5-HT₄ gene. All other examined tissues showed bands corresponding to either the a or b splice variant, but not to the h variant.

Transient expression of the 5-HT_{4(h)} variant in mammalian cells and pharmacological characterization

In order to compare the pharmacological properties of the 5-HT_{4(h)} variant with those of the previously cloned 5-HT_{4(a)} and b variant, the three corresponding pcDNA3 expression constructs were transiently transfected into COS cells. The receptors were investigated by radioligand binding assays on membrane preparations. Saturation analysis experiments were performed with the agonist [³H]5-HT as well as with the antagonist [³H]GR113808. Antagonist and agonist saturation binding of the h variant from 3 independent transfections resulted in a straight line in Scatchard analysis, revealing a single high-affinity binding site. The ligand concentration isotherms of the h 5-HT_{4(h)} revealed a K_D of 0.24 (+/-0.17) nM using the antagonistic ligand and 1.65 (+/-0.55) nM based on the agonistic ligand (Fig. 3A and B).

These values were not significantly different from the two other variants we investigated as reference, the K_D of the variants a and b were respectively 0.14 (+/-0.06) and 0.19 (+/-0.07) nM for [³H]GR113808

- 33 -

saturation binding and 3.7 (± 0.6) and 4.6 (± 1.9) nM for [3 H]5-HT saturation binding. The h 5-HT_{4(h)} receptor displayed a B_{max} of 605 (± 174) fmol/mg of protein with [3 H]GR113808, and 61 (± 12) fmol/mg of protein with [3 H]5-HT. Scatchard analysis from [3 H]5-HT binding suggested two affinity sites for the two 5-HT₄ receptor variants a and b. For each of the 3 independent transfections, COS-7 cells transfected with the vector alone showed no specific binding using the radioligand [3 H]GR113808. Saturation binding experiments with [3 H]GR113808 and [3 H]5-HT were performed in parallel for each of the 3 independent transfections. The resulting ratio of the B_{max} found using the antagonistic radioligand versus the B_{max} based on the agonistic radioligand reveals the proportion of the coupled receptors among the total number of receptors. We found this proportion to be 9.8 (± 1.05) for the h5-HT_{4(h)}. The same result was obtained for the B_{max} values found for the high affinity sites of the 2 reference 5-HT receptor variants a and b. However using the B_{max} values obtained with the low affinity site, yielded a ratio of coupled receptors to total receptors that was 3 times lower. This result suggests a difference in the G-protein coupling of the h 5-HT_{4(h)} variant, compared to the h 5-HT_{4(a)} and h 5-HT_{4(b)} variant. The pharmacological binding profile of the h5-HT_{4(h)} was studied by competition binding assays using six different agonists and two antagonists in combination with the radioligand [3 H]GR113808 at a concentration of 0.25 nM (Fig. 4) on COS-7 membranes. Results are the mean of three independent transfections.

- 34 -

The 5-HT₄ receptor is suggested to be involved in a number of different physiological processes, which makes it an important pharmacological target. 5-HT₄ receptor activation influences gastrointestinal motility (Meulemans and Schuurkes, 1992), bladder function (Candura et al., 1996), exerts chronotropic and inotropic effects at the heart (Kaumann et al., 1990) and centrally enhances striatal dopamine release (Bonhomme et al., 1995) as well as associative memory in rats (Marchetti-Gauthier et al., 1997). This variety of physiological effects is paralleled by a variety of splice variants which have been discovered in the course of the last two years (Gerald et al., 1995; Claeyssen et al., 1996; Van den Wyngaert et al., 1997; Claeyssen et al., 1997; Blondel et al., 1997; Blondel et al., 1998). For these splice variants up to now no well documented specific biochemical or physiological properties described. However for splice variants of other GPCRs more information is available. For the C-terminal splice variants of the prostaglandin EP₃ receptor coupling to different G-proteins and signal transduction systems has been shown (Namba et al., 1993). C-terminal splice variants of the mouse somatostatin receptor differ in the efficiency of adenylate cyclase inhibition and receptor desensitization (Vanetti et al., 1993). Differential splicing at the third intracytoplasmic loop of the PACAP receptor leads to coupling to different G- proteins (Spengler et al., 1993), the same was found for C-terminal splice variants of mGluR1 (Pin et al. 1992). Pickering et al. (1993) showed for the same variants in addition differential intracellular distribution. For some of these GPCR

- 35 -

splice variants also differential tissue distribution has been shown (Spengler et al., 1993), also this is not the case for others (Pin et al., 1992). The up to now published 5-HT₄ splice variants all vary in their C-terminus, which suggests in analogy to the results obtained from other GPCRs that they may differ from each other in respect to G-protein usage, desensitization and/or subcellular localization. The 5-HT_{4(h)} variant described in this study has an extra insertion of 14 amino acids in the second extracellular loop, to our knowledge there are no other descriptions of such a variation produced by alternative splicing. Surprisingly this modification led to the loss of the low affinity agonist binding site that was found for the a and b splice variant. Although the ratio of coupled to uncoupled receptor is still the same for the high affinity agonist binding site across all three known variants, it is likely that the insertion of 14 amino acids into the second extracellular loop leads to a change of receptor topology that is reflected in G-protein binding. This change in receptor topology is also suggested by the agonistic effect that GR113808 exerts at this variant, which acts as an antagonist at all other 5-HT₄ variants. The availability of a variety of 5-HT₄ splice variants offers the opportunity for medicinal chemistry to pursue a higher degree of specificity for drug development. Given that the standard 5-HT₄ receptor antagonist GR 113 808 showed agonistic activity on the 5-HT_{4(h)} receptor variant and that 5-HT₄ receptor antagonists are under investigation as compounds for the treatment of irritable bowel syndrome (IBS), testing of any given ligand on the 5-

- 36 -

HT4(h) variant is essential before classifying it as an 5-HT4 receptor antagonist. Based on the specific tissue distribution of the 5-HT_{4(h)} to the LES, compounds showing specificity towards that 5-HT₄ splice variant may have therapeutic value for the treatment of heartburn, reflux, irritable bowel syndrome, esophagitis, Barrett's esophagus, esophageal cancer, achalasia, esophageal stenosis, esophageal spasms, esophageal hiatal hernia or other esophageal motility disorders. Furthermore these compounds may be of value in the treatment of airway disorders possibly connected with oesophageal irritation, such as asthma, bronchospasms, aspiration and its consequences (bronchitis, (broncho)pneumonia, bronchiectasia).

Compounds acting on 5-HT₄ receptor splice variants found in the lower oesophageal sphincter, may additionally be useful in treating or alleviating the symptoms of diseases of the lower oesophageal sphincter or other conditions such as achalasia; oesophageal stenosis (due to systemic sclerosis, tumours, burns) or compression, oesophageal spasms or other oesophageal motility disorders, irritable bowel syndrome, asthma, bronchospasms and other airway disorders possibly connected with oesophageal irritation aspiration and its consequence (bronchitis, (broncho)pneumonia, bronchiectasia, ...); (hiatus) hernia; denervation of the oesophagus (e.g. after certain types of trauma or surgery), disturbances in oesophageal innervation; pregnancy (not a disease or even a condition that as such could be treated with 5HT4-receptor compounds, but one in which -for various reasons-oesophageal reflux and its consequences are

- 37 -

more common); emesis; postoperative ileus; diabetic gastroparesis.

Abbreviations used: AC, adenylyl cyclase; DMEM,
5 Dulbecco's modified Eagle medium; DMSO, dimethyl
sulfoxide; 5-HT, 5-hydroxytryptamine, serotonin; Ki,
inhibition constant; LES, lower esophageal sphincter;
ORF, open reading frame.

10

References

- Blondel, O., Vandecasteele, G., Gastineau, M.,
Leclerc, S., Dahmoune, Y., and Fischmeister, R.
5 (1997) Molecular and functional characterization
of a 5-HT₄ receptor cloned from human atrium. *FEBS
Letters* **412**, 465-474.
- Blondel, O., Gastineau, M., Dahmoune, Y., Langlois,
M., and Fischmeister, R. (1998) Cloning,
10 expression, and pharmacology of four human 5-
hydroxytryptamine₄ receptor isoforms produced by
alternative splicing in the carboxyl terminus. *J.
of Neurochemistry* **70**, 2252-2261.
- Bonhomme, N., De Deurwaerdere, P., Le Moal, M., and
15 Spampinato, U. (1995) Evidence for 5-HT₄ receptor
subtype involvement in the enhancement of striatal
dopamine release induced by serotonin: a
microdialysis study in the halothane-anesthetized
rat. *Neuropharmacology* **34(3)**, 269-279.
- 20 Buchheit, K.-H. and Buhl, T. (1991) Prokinetic
benzamides stimulate peristaltic activity in the
isolated guinea pig ileum by activation of 5-HT₄
receptors. *Eur. J. Pharmacol.* **205**, 203-208.
- Candura, S.M., Messori, E., Franceschetti, G.P.,
25 D'Agostino, G., Vicini, D., Tagliani M., and
Tonini M. (1996) Neural 5-HT₄ receptors in the
human isolated detrusor muscle: effects of indole,
benzimidazolone and substituted benzamide agonists
and antagonists. *Br. J. Pharmacol.* **118**, 1965-1970.
- 30 Claeysen, S., Sebben, M., Laurent, J., Bockaert, J.,
and Dumuis, A. (1996) Cloning, expression and
pharmacology of the mouse 5-HT_{4L} receptor. *FEBS*

Letters 398, 19-25.

- 5 Claeyssen, S., Faye, P., Sebben, M., Lemaire, S.,
Bockaert, J., and Dumuis, A. (1997) Cloning and
expression of human 5-HT₄S receptors. Effect of
receptor density on their coupling to adenylyl
cyclase. *NeuroReport* 8, 3189-3196.
- 10 Dumuis, A., Bouhelal, R., Sebben, M., Cory, R., and
Bockaert, J. (1988) A non-classical 5-
hydroxytryptamine receptor positively coupled with
adenylate cyclase in the central nervous system.
Mol. Pharmacol. 34, 880-887.
- Eglen, R.M., Wong, E.H.F., Dumuis, A., and Bockaert,
J. (1995) Central 5-HT₄ receptors. *Trends in
Pharmacol. Sci.* 16, 391- 398.
- 15 Elswood, C.J., Bunce, K.T., and Humphrey, P.P.A.
(1991) Identification of putative 5-HT₄ receptors
in guinea-pig ascending colon. *Eur. J.
Pharmacology* 196, 149-155.
- 20 Fontana D.J., Daniels, S.E., Wong, E.H., Clark, R.D.
and Eglen, R.M. (1997) The effects of novel 5-
hydroxytryptamine (5-HT)₄ receptor ligands in rat
spatial navigation. *Neuropharmacology* 36 (4), 689
- 696.
- 25 Galeotti, N., Ghelardini, C., and Bautolini, A. (1998)
Role of 5-HT₄ receptors in the mouse passive
avoidance test. *J. Pharmacol. & Exp. Therapeutics*
286 (3), 1115-1121.
- 30 Gerald, C., Adham, N., Kao, H.-T., Olson, M.A., Laz,
T.M., Schechter, L.E., Bard, J.A., Vaysse, J.-J.,
Hartig P.R., Branchek, T.A., and Weinshank, R.L.
(1995) The 5-HT₄ receptor: molecular cloning and
pharmacological characterization of two splice

- 40 -

- variants. *EMBO J.* **14** (12), 2806-2815.
- Hedge, S.S. and Eglén, R.M. (1996) Peripheral 5-HT₄ receptors. *FASEB J.*, **10** (12), 1398 -1407.
- 5 Kaumann, A.J., Sanders, L., Brown, A.M., Murray, K.J. and Brown, M.J. (1990) A 5-hydroxytryptamine receptor in human atrium. *Br. J. Pharmacol.* **100**, 879-885.
- 10 Lefebvre, H., Contesse, V., Delarue, C., Fevilloley, M., Héry, F., Grise, P., Raynaud, G., Verhofstad, A.A.J., Wolfe, L.M. and Vaudry, H. (1992) *Neuroscience* **47**, 999
- Maniatis, T., Fritsch, E.F., and Sambrook, J. (1982) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor, New York.
- 15 Marchetti-Gauthier, E., Roman, F.S., Dumuis, A., Bockaert, J., and Soumireu-Mourat, B. (1997) BIMU1 increases associative memory in rats by activating 5-HT₄ receptors. *Neuropharmacology* **36** (4/5), 697-706.
- 20 Meulemans, A.L. and Schuurkes, A.J. (1992) Is the action of cisapride on the guinea pig ileum mediated via 5-HT₄ receptors. *Eur. J. of Pharmacol.* **212**, 51-59.
- 25 Monferini, E., Gaetani, P., Baena, R. R., Giraldo, E., Parenti, M., Zocchetti, A., and Rizzi, C.A. (1993) Pharmacological characterization of the 5-Hydroxytryptamine receptor coupled to adenylyl cyclase stimulation in human brain. *Life Sciences* **52**, 61-65.
- 30 Moummi, C., Yang, D.-C., and Gullikson, G. W. (1992) 5-HT₄ receptor activation induces relaxation and associated cAMP generation in rat esophagus. *Eur.*

J. Pharmacol. **216**, 47-52.

- Namba, T., Sugimoto, Y., Negishi, M., Irie, A.,
Ushikubi, F., Kakizuka, A., Ito, S., Ichikawa, A.,
and Narumiya, S. (1993) Alternative splicing of C-
5 terminal tail of prostaglandin E receptor subtype
EP3 determines G-protein specificity. *Nature* **365**,
166-170.
- Pickering, D.S., Thomsen, C., Suzdak, P.D., Fletcher,
E.J., Robitaille, R., Salter, M.W., MacDonald,
10 J.F., Huang, X.-P., and Hampson, D.R. (1993) A
comparison of two alternatively spliced forms of a
metabotropic glutamate receptor coupled to
phosphoinositide turnover. *J. of Neurochemistry*
61, 85-92.
- 15 Pin, J.-P., Waeber, C., Prezeau, L., Bockaert, J., and
Heinemann, S.H. (1992) Alternative splicing
generates metabotropic glutamate receptors
inducing different patterns of calcium release in
Xenopus oocytes. *Proc. Natl. Acad. Sci. U.S.A.* **89**,
20 10331-10335.
- Silvestre, J.S., Fernandez, A.J., and Palacios, J.M.
(1996) Effects of 5-HT₄ receptor antagonists on
rat behaviour in the elevated plus - maze test.
Eur. J. Pharmacol. **309(3)**, 219-222.
- 25 Spengler, D., Waeber, C., Pantaloni, C., Holsboer, F.,
Bockaert, J., Seeburg, P., and Journot, L. (1993)
Differential signal transduction by five splice
variants of the PACAP receptor. *Nature* **365**, 170-
175.
- 30 Ullmer, C., Schmuck, K., Kalkman, H.O., and Lübbert,
H. (1995) Expression of serotonin receptor mRNAs
in blood vessels. *FEBS Letters* **370**, 215-221.

- Van den Wyngaert, I., Gommeren, W., Verhasselt, P.,
Jurzak, M., Leysen, J., Luyten, W. and Bender, E.
(1997) Cloning and expression of a human serotonin
5-HT₄ receptor cDNA. *J. of Neurochemistry* **69**,
5 1810-1819.
- Vanetti, M., Vogt, G., and Höllt, V. (1993) The two
isoforms of the mouse somatostatin receptor
(mSSTR2A and mSSTR2B) differ in coupling
efficiency to adenylate cyclase and in agonist-
10 induced receptor desensitization. *FEBS Letters* **331**
(3), 260-266.
- Vilaro, M.T., Cortes, R., Gerald, C., Branchek, T.A.,
Palacios, J.M., and Mengod, G. (1996) Localization
of 5-HT₄ receptor mRNA in rat brain by in situ
15 hybridization histochemistry. *Mol. Brain Res.* **43**,
356-360.

Table 1 Comparison of potency of compounds to compete with 0.25 nM [³H]GR113808 binding in membranes from COS-7 cells transiently transfected with respectively human 5-HT_{4(h)}, 5-HT_{4(a)}, or 5-HT_{4(b)} receptors.

Compounds	Human 5-HT ₄ h receptor in Cos-7 cells		Human 5-HT ₄ a receptor in Cos-7 cells		Human 5-HT ₄ b receptor in Cos-7 cells	
	pIC50 +/- SD (n)	pK _i	pIC50 +/- SD (n)	pK _i	pIC50 +/- SD (n)	pK _i
Agonist						
5HT	6.09 +/- 0.05 (3)	6.43	6.80 +/- 0.37 (3)	6.89	6.47 +/- 0.18 (3)	6.63
Prucalopride	6.74 +/- 0.24 (3)	8.03	7.20 +/- 0.40 (4)	7.21	7.04 +/- 0.25 (4)	7.16
Cisapride	6.78 +/- 0.32 (3)	7.05	7.22 +/- 0.47 (4)	7.23	6.92 +/- 0.33 (4)	6.99
Mosapride	6.55 +/- 0.24 (3)	6.85	6.48 +/- 0.19 (3)	6.71	6.04 +/- 0.27 (3)	6.18
SDZ-HTF919	7.45 +/- 0.52 (3)	7.59	8.08 +/- 0.66 (4)	7.86	7.89 +/- 0.20 (4)	8.09
5-MeOT	5.82 +/- 0.27 (3)	6.1	6.09 +/- 0.53 (4)	6.08	6.11 +/- 0.52 (4)	6.13
Antagonist						
SB204070	10.01 +/- 0.17 (3)	10.38	10.07 +/- 0.32 (4)	10.1	9.98 +/- 0.16 (4)	10.04
GR113808	9.05 +/- 0.31 (3)	9.29	9.45 +/- 0.35 (4)	9.54	9.39 +/- 0.13 (4)	9.57

Table 2 Comparison of potency and affinity of agonist compounds to stimulate cAMP formation in COS-7 cells transiently transfected with respectively human 5-HT_{4(h)}, 5-HT_{4(a)} or 5-HT_{4(b)} receptors.

% of 5HT max (+/-SD)				pEC50 (+/1 SD)			
5-HT4 variants		a	b	h	a	b	h
Compounds		n=5	n=8	n=3	n=5	n=8	n=3
10	5HT	100	100	100	8.24	8.31	7.9
		-	-	-	(+/-0.156)	(+/-0.147)	(+/-0.26)
	Prucalopride	118	117.36	95	8.24	8.49	8.36
15		(+/-7.2)	(+/-3.4)	(+/-11.6)	(+/-0.105)	(+/-0.049)	(+/-0.62)
	Cisapride	98.17	113.62	123.53	7.82	7.9	7.99
		(+/-7.5)	(+/-4.36)	(+/-20.85)	(+/-0.26)	(+/-0.113)	(+/-0.62)
	GR113808	0	0	54.6	-	-	7.95
				(+/-5.8)			(+/-0.38)

Claims

1. An isolated or substantially pure form of a
nucleic acid molecule encoding a human 5-HT_{4(h)}
5 receptor.
2. The nucleic acid molecule of claim 1 encoding a
human 5-HT_{4(h)} receptor comprising the amino acid
sequence illustrated in Figure 1b or encoding a
10 functional equivalent, derivative or bioprecursor
of said receptor.
3. A nucleic acid molecule according to claim 1 or 2
which is a DNA molecule.
15
4. A nucleic acid molecule according to claim 3,
wherein said DNA molecule is a cDNA molecule.
5. A nucleic acid molecule according to any of
20 claims 2 to 4 comprising the sequence illustrated
in Figure 1b.
6. A nucleic acid molecule capable of hybridising to
the molecule of any of claims 1 to 5 or the
25 complementary sequences thereto under conditions
of high stringency.
7. A human 5-HT_{4(h)} receptor encoded by the nucleic
acid molecule according to any of claims 1 to 5.
30
8. A DNA expression vector comprising a nucleic acid
molecule according to any of claims 3 to 5.

9. A host cell transformed or transfected with the vector of claim 8.
10. A host cell according to claim 9, which cell is a mammalian cell.
11. A host cell according to claim 10, which mammalian cell is a COS-7 cell.
12. A transgenic cell, tissue or organism comprising a transgene capable of expressing a human 5-HT_{4(h)} receptor protein comprising the amino acid sequence of Figure 1b or an amino acid sequence of a functional equivalent, derivative or bioprecursor of said receptor.
13. A transgenic cell, tissue or organism according to claim 12 wherein said transgene comprises a nucleic acid molecule according to any of claims 1 to 5.
14. A human 5-HT_{4(h)} receptor protein or a functional equivalent, derivative or bioprecursor thereof, expressed by the cell according to any of claims 9 to 11 or the cell tissue or organism according to claim 12.
15. A HEK 293 or COS-7 5-HT_{4(h)} cell line transfected with the expression vector of claim 8.
16. An antisense molecule comprising a nucleic acid molecule which is capable of hybridising to the nucleic acid of any of claims 1 to 5 under

conditions of high stringency.

- 5 17. A pharmaceutical composition comprising a molecule according to claim 16 together with a pharmaceutically acceptable carrier, diluent or excipient therefor.
- 10 18. An antisense molecule according to claim 13 for use as a medicament.
- 15 19. A purified or isolated human 5-HT_{4(h)} receptor protein comprising the amino acid sequence according to Figure 1b or the amino acid sequence of a functional equivalent, derivative, fragment or bioprecursor of said sequence.
- 20 20. A pharmaceutical composition comprising a molecule according to any of claims 1 to 5 together with a pharmaceutically acceptable carrier, diluent or excipient therefor.
- 25 21. An antagonist or an agonist of a ligand of the human 5-HT_{4(h)} receptor protein according to any of claims 14 or 19.
- 30 22. A pharmaceutical composition comprising an antagonist or an agonist according to claim 21 together with a pharmaceutically acceptable carrier, diluent or excipient therefor.
23. A method of determining whether a compound is an agonist or an antagonist of a ligand of a human 5-HT_{4(h)} receptor, which method comprises

contacting a cell according to any of claims 7 to 10 expressing said receptor protein with said compound in the presence of said ligand and monitoring cAMP formation in said cell.

5

24. A method according to claim 23 wherein said cell is a human cell.

10

25. A method of determining whether a compound binds to a human 5-HT_{4(h)} receptor which method comprises contacting a cell, according to any of claims 9 to 12 or a membrane preparation comprising said receptor, with said compound and establishing the binding affinity of said compound for said receptor.

15

26. A compound identifiable as an agonist or antagonist according to the method of claim 24 or 25.

20

27. A compound according to claim 26 for use as a medicament.

25

28. Use of a compound identifiable according to the method of claim 26 or an antisense molecule according to claim 16 in the manufacture of a medicament for the treatment of any of heartburn, reflux, esophagitis, Barrett's esophagus, esophageal cancer, achalasia, esophageal stenosis, esophageal spasms, esophageal hiatal hernia or other esophageal motility disorders, oesophageal irritation, such as asthma, bronchospasms, aspiration and its consequences

30

(bronchitis, (broncho)pneumonia, bronchiectasia) and other diseases of the lower oesophageal sphincter, or achalasia; oesophageal stenosis (due to systemic sclerosis, tumours, burns, or the like) or compression, oesophageal spasms or other oesophageal motility disorders, asthma, irritable bowel syndrome, bronchospasms and other airway disorders possibly connected with oesophageal irritation aspiration and its consequence (bronchitis, (broncho)pneumonia, bronchiectasia); (hiatus) hernia; denervation of the oesophagus (e.g. after certain types of trauma or surgery), disturbances in oesophageal innervation.

29. A pharmaceutical composition comprising a compound according to claim 27 together with a pharmaceutically acceptable carrier diluent or excipient therefor.

30. An antibody specific for a human 5-HT_{4(h)} receptor according to claim 7 or 19.

31. A kit for determining whether a compound is an agonist or an antagonist of a 5-HT_{4(h)} ligand, which kit comprises a cell according to any of claims 9 to 12, means for contacting said compound and said ligand with said cell and means for measuring cAMP formation in said cell.

32. A kit according to claim 31 wherein said cell is a COS-7 cell.

33. A pharmaceutical composition incorporating the nucleic acid sequence according to any of claims 1 to 5, or the antibody according to claim 30, together with a pharmaceutically acceptable carrier, diluent or excipient therefor.

34. A method of identifying a ligand for 5-HT_{4(h)} receptor, which method comprises contacting a cell expressing said receptor with said compound to be tested and monitoring the level of any 5-HT_{4(h)} mediated functional or biological response.

FIG. 1A

FW ON start

h5HT4B	: GTAATGGACAAACTTGATGCTAATGCTGAGTTCTGAGGAGGGTTTCGGGTCAAGTGAGAGGTTGGTGTCTCGACGGTTATCTGATGGCCATCTTGGGGAACCTG	: 117
d5HT4B	: ---ATGGACAAACTTGATGCTAATGCTGAGTTCTGAGGAGGGTTTCGGGTCAAGTGAGAGGTTGGTGTCTCGACGGTTATCTGATGGCCATCTTGGGGAACCTG	: 114
h5HT4B	: CTGGTGATGGTGGCTGTGCTGGGACAGGACGCTCAGGAAATATAAAACAAATATTTCATTGTATCTCTTGTCTTTGGGGATCTGCTGGTTTCGGTGCTGGTGATGCCCTTTGGT	: 234
d5HT4B	: CTGGTGATGGTGGCTGTGCTGGGACAGGACGCTCAGGAAATATAAAACAAATATTTCATTGTATCTCTTGTCTTTGGGGATCTGCTGGTTTCGGTGCTGGTGATGCCCTTTGGT	: 231
h5HT4B	: GCCATTGAGCTGGTTCAAGACATCTGGATTATAGGGAGGTTGTTGTCTGTTTCGGACATCTCTGGACGTCCTGCTCAACACGGCATCGATTTTTCACCTGTGCTGCATTTCTCTG	: 351
d5HT4B	: GCCATTGAGCTGGTTCAAGACATCTGGATTATAGGGAGGTTGTTGTCTGTTTCGGACATCTCTGGACGTCCTGCTCAACACGGCATCGATTTTTCACCTGTGCTGCATTTCTCTG	: 348
h5HT4B	: GATAGGTATTAGGCCATCTGCTGCCAGGCTTTGGTCTATAGGAACAAGATGACCCCTCTGGCATCGCATTAATGCTGGAGGCTGCTGGGTCAATCCACAGTTTATTTCTTTCTC	: 468
d5HT4B	: GACAGGTATTATGCCATCTGCTGCCAGGCTCTGGTCTATAGGAACAAGATGACCCCTCTGGCATCGGCTTAATGCTGGAGGCTGCTGGATCATCCCATGTTTATCTCTTTCTC	: 465
h5HT4B	: CCTATAATGCAAGGCTGGAATAACATTGGCATAATTGATTGGAAGGAGTCTAAACCAAGGCTGGGCCAGGATTTTCATGCCATAGAAAAGAGGAGTTCAACCAAGAACTCTTAAC	: 585
d5HT4B	: CCTATAATGCAAGGCTGGAATAATATTGGCATAATTGATTGGAAGGAGTATCCAAAGGCTGGGCCAGGATTTGCAATGTCATAGAAAAGAGGAGTTCAACCAAACTCCAAAC	: 582
h5HT4B	: TCTACGTACTGTGCTTTCATGGTCAACAGCCCTACGCCATCACCTGCTGTGGTGGGCTTCTACATCCCATTTCTCTCATGGTGTGGGCTATTACCGCATCTATGTCACAGCT	: 702
d5HT4B	: TCTACGTACTGTATCTTTCATGGTCAACAGCCCTACGCCATCACCTGCTGTGGTGGGCTTCTACATCCCATTTCTCTCATGGTGTGGGCTATTATGTCATCTACGTACAGCA	: 699

1 / 8

FW AB1

FW B1

FW AB3

REV B2

2 / 8

		FW AB2	
h5HT4B	:	AAGGAGCATGCCCATCCAGATCCAGATGTTACAGGGCAGGAGCCCTCTCCGAGAGCAGGCCCTCAGTGGGACAGCCAGCATAGCACTCATCGCATGAGGACACAGACCAAGCAGCC	: 819
d5HT4B	:	AAAGAGCATGCCCATCCAGATCCAGATGTTACAGGGCAGGAGCCCTCTGAAAGGCAGGCCAGCTGACCCAGCAGCACTCATCGCATGAGGACCCGAGACCAAGCAGCC	: 816
		REV AB2	
h5HT4B	:	AAGACCTGTGCAATCATATGGGTTGCTTCTGCTGCTGGGACCAATCTTTGTCAACCAATATTGTGGATCCCTTTTCATAGACTACACTGTCCCTGGGACGGTGTGGACTGCTTTC	: 936
d5HT4B	:	AAGACCTGTGCAATCATATGGGTTGCTTCTGCTGCTGGGCTCCATCTTGTCTACTAATATTGTGGATCCATTCATAGACTACACTGTCTGCTGGGACGGTGTGGACTGCTTTC	: 933
h5HT4B	:	CTCTGGCTGGGCTATATCAATTCCGGGTTGAACCCCTTTTCTCTAGCCCTTCTTTGAATAAGTCTTTTAGAGGTGCCCTTCCTCATCATCTCTGCTGTGATGATGAGCCCTACCGAAGA	: 1053
d5HT4B	:	CTCTGGCTGGGCTACATCAACTCTGGGTTGAACCCCTTTCTGTAGCCCTTCTTTGAATAAGTCTTTTAGAGGTGCCCTTCCTCATCATCTCTGCTGTGATGATGAGCCCTACCGAAGA	: 1050
h5HT4B	:	CCTTCCATTCTGGGCCAGACTGTCCCTTGTTCACCCACAAACCATTAAATGGATCCACACATGTACTAAGGATGCAAGTGTGGTGTGGCCAGTGGGAGAGTCAGTGTACCCCGCCA	: 1170
d5HT4B	:	CCTTCCATTCTGGGCCAGACTGTCCCTTGTTCACCCACAAACCATTAAATGGATCCACACATGTACTAAGGATGCAAGTGTGGTGTGGCCAGTGGGAGAGTCAGTGTACCCCGCCA	: 1167
		REV LOI	
h5HT4B	:	GCAACTTCTCTTTTGGTGGTGTGCTCAGCCCAAGTGCACACTTAGGCCCTGGGACAATGACCCAGAACAGACAGCCCATGCCCTCCGAAAGAGGGGCCAGGTCTTAAGTGTGCTTTC	: 1281
d5HT4B	:	GCAACTTCTCTTTTGGTGGTGTGCTCAGCCCAAGTGCACACTTAG-----	: 1209
		REV 3non	

FIG. 1Acont'd

3 / 8

h5HT4B	:	VMDKLDANVSSEEGFGSVEKVLLTFLSTVILMAILGNLLVMVAVCWDRQLRKIKITNYFIVSLAEADLLSVLVMPE	:	77
d5HT4B	:	-MDKLDANVSSEEGFGSVEKVLLTFLSTVILMAILGNLLVMVAVCWDRQLRKIKITNYFIVSLAEADLLSVLVMPE	:	76
h5HT4B	:	GAIELVQDIWIYGEVFCIVRTSLDVLLTTASIFHLCCISLDRYYAICCPPLVYRNKMTPLRIALMLGGCWVPTFIS	:	154
d5HT4B	:	GAIELVQDIWIYGEVFCIVRTSLDVLLTTASIFHLCCISLDRYYAICCPPLVYRNKMTPLRIALMLGGCWVPTFIS	:	153
h5HT4B	:	FLPIMQGWNNIGIIDLERSLNQGLGQDFHAEKRRKFNQNSNSTYCVFMVKNKPYAITCSVVAFYIPELLMVLAYYRIY	:	231
d5HT4B	:	FLPIMQGWNNIGIIDLERISKPRLGQDLHVIEKRRKFNQNSNSTYCIEMVKNKPYAITCSVVAFYIPELLMVLAYYRIY	:	230
h5HT4B	:	VTAKEHAHQIQMLQRAGASSESRPOSADQHSRMRRTETKAAKTLICIIMGCFCLCWAPFEFTNIVDPFIDYTVPGQV	:	308
d5HT4B	:	VTAKEHAHQIQMLQRAGAPSEGRPPADQHSRMRRTETKAAKTLICIIMGCFCLCWAPFEFTNIVDPFIDYTVPGQV	:	307
h5HT4B	:	WTAFLWLGYINSGLNPFYAFLNKSFERRAFLLIILCCDDERYRRRPSILGQTVPCSTTTINGSTHVLDAVECGGQWES	:	385
d5HT4B	:	WTAFLWLGYINSGLNPFYAFLNKSFERRAFLLIILCCDDERYRRRPSILGQTVPCSTTTINGSTHVLDAVECGGQWES	:	384
h5HT4B	:	QCHPPATSPLVAAQPSDTAPGTMTQKTAMPPKEGQVLSCL	:	426
d5HT4B	:	QCHPPATSPLVAAQPSDT-----	:	402

FIG. 1B




FIG. 2

5 / 8




FIG. 3

6 / 8




FIG. 4

7 / 8

Stimulation of AC through human 5-HT₄h receptor in COS-7 cells




FIG. 5

8 / 8




FIG. 6

Description of Sequences

- 5 SEQ ID NO. 1 corresponds to the nucleic acid
 molecule encoding 5-HT_{4(h)} receptor
 illustrated in Figure 1a.
- SEQ ID NO. 2 corresponds to the amino acid sequence
 of 5-HT_{4(h)} receptor illustrated in
 Figure 1b.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/05592

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N5/10 C07K14/705 C07K16/28 A61K38/17
G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBL, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	BENDER E. ET AL.: "Structure of the human serotonin H-HT4 receptor gene and cloning of a novel 5-HT4 splice variant." J. NEUROCHEM., vol. 74, February 2000 (2000-02), pages 478-489, XP000953119 the whole document	1-20, 23-25, 28,30-34
A	BLONDEL O. ET AL.: "Cloning, expression and pharmacology of four human 5-hydroxytryptamine-4 receptor isoforms produced by alternative splicing in the carboxy terminus." J. NEUROCHEM., vol. 70, June 1998 (1998-06), pages 2252-2261, XP000953121 the whole document	1-20, 23-25, 28,30-34
	--- -/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

25 October 2000

Date of mailing of the international search report

07/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/EP 00/05592

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 94 14957 A (GERALD CHRISTOPHE ;WEINSHANK RICHARD L (US); HARTIG PAUL (US); BRA) 7 July 1994 (1994-07-07) abstract claims 1-93</p>	<p>1-20, 23-25, 28, 30-34</p>

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 21,22,26,27 and partly 28

Claims 21,22,26,27 and partly 28 refer to agonists and antagonists of the new 5-HT₄ splice variant protein, without however giving a true technical characterization. Moreover, no such compounds are defined in the application, and the agonists and antagonists known to act on other splice forms do not necessarily have the same effect on the present splice form (Bender, J. Neurochem. 74:478-489, 2000).

In consequence, said claims are ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported in the description pursuant to Art. 5 and 6 PCT. No search can be carried out for such purely speculative claims, the wording of which is, in fact, a mere recitation of the results to be achieved.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Patent Application No

PCT/EP 00/05592

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9414957 A	07-07-1994	US 5472866 A	05-12-1995
		AU 685076 B	15-01-1998
		AU 5960694 A	19-07-1994
		CA 2129969 A	07-07-1994
		EP 0642578 A	15-03-1995
		US 5766879 A	16-06-1998